

Catalogue Number	Product	Order number / Unit
101	Mannose Triflate, ultra pure Precursor for [¹⁸F]FDG (2-[¹⁸F]Fluoro-2-deoxy-D-glucose) packing for GE TRACERlab MXFDG Molar Mass: 480.37 C ₁₅ H ₁₉ F ₃ O ₁₂ S [92051-23-5] Colourless or nearly colourless crystals packaged in clear glass vials (10 ml headspace) with teflon-faced rubber stoppers, center tear-off crimp caps, argon flushed. Melting range 119 - 122 degC. Soluble in acetonitrile, acetone, DMSO, methanol. Insoluble in aqueous media. Purity: > 99 % Certificates: CoA; ¹ H and ¹⁹ F NMR spectra; HPLC; IR spectrum; melting point Chemical Name: CA index name: beta-D-mannopyranose, 1,3,4,6-tetraacetate 2-(trifluoromethanesulfonate) Synonymes: TATM; mannose triflate; 1,3,4,6-tetra-O-acetyl-2-O-trifluoro-methanesulfonyl-beta-D-mannopyranose Literature: Hamacher K. et al. Efficient stereospecific synthesis of no-carrier-added 2-[¹⁸ F]fluoro-2-deoxy-D-glucose using aminopolyether supported nucleophilic substitution. J. Nucl. Med. 1986, 27, 235-238. Padgett H. et al. Computer-controlled radiochemical synthesis: a chemistry process control unit for the automated production of radiochemicals. Appl. Radiat. Isot. 1989, 40, 433-445. Pavliak V. et al. A short synthesis of 1,3,4,6-Tetra-O-acetyl-2-azido-2-deoxy-beta-D-glucopyranose and the corresponding α-glucosyl chloride from D-mannose. Carbohydr. Res. 1991, 210, 333-337. Chirakal R. Traces of fluorine containing impurities in the mannose triflate and their adverse effect on the radiochemical yield of 2-[¹⁸ F]FDG. XIIth ISRC. Uppsala, Sweden 1997, 214-216.	101.0020: 20 mg per vial 101.0025: 25 mg per vial 101.0040: 40 mg per vial 101.0062: 62 mg per vial Please inquire for customized filling and bulk quantities.

